

## 2013 Appraisal Index

To assist clinicians looking for authoritative assistance with clinical problems, the journal publishes an annual index of content from the most recent two years of Appraisal pages. This index includes content from Volumes 58 and 59 of *Journal of Physiotherapy*. Content is indexed under the PEDro codes: subdiscipline, intervention, problem, and body part. It is identified by Appraisal section and Volume and page number. Some content is indexed under more than one code.

### SUBDISCIPLINE

#### Cardiothoracics

##### CAPs

- Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD) Vol 58 No 1, p 60
- Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease Vol 58 No 2, p 129
- Exercise training improves cardiovascular fitness in people receiving haemodialysis for chronic renal disease Vol 58 No 2, p 130
- Interval training confers greater gains than continuous training in people with heart failure Vol 58 No 3, p 199
- Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease Vol 58 No 4, p 270
- Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke Vol 58 No 4, p 271
- Long-term aerobic exercise maintains peak VO<sub>2</sub>, improves quality of life, and reduces hospitalisations and mortality in patients with heart failure Vol 59 No 1, p 56
- Exercise training following lung transplant is now evidence-based practice Vol 59 No 1, p 58
- Telehealth reduces hospital admission rates in patients with COPD Vol 59 No 2, p 129
- Sun-style T'ai Chi improves walking endurance and health-related quality of life in people with COPD Vol 59 No 4, p 274

##### Clinical Practice Guidelines

- Paediatric neuromuscular weakness Vol 59 No 2, p 133
- Cardiovascular disease risk Vol 59 No 2, p 133
- Chronic obstructive lung disease Vol 59 No 4, p zzz

#### Continence & Women's Health

##### CAPs

- A 12-week exercise program performed during the second trimester does not prevent gestational diabetes in healthy pregnant women Vol 58 No 3, p 198

##### Clinimetrics

- Active Straight Leg Raise Vol 58 No 2, p 132

#### Ergonomics & Occupational Health

##### Clinimetrics

- The Work Limitations Questionnaire (WLQ-25) Vol 58 No 4, p 277

##### Clinical Practice Guidelines

- The Work Instability Scale Vol 59 No 3, p 212

#### Gerontology

##### CAPs

- Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility Vol 58 No4, p 269
- Tai Chi training is effective in reducing balance impairments and falls in patients with Parkinson's disease Vol 59 No 1, p 55
- Body-weight supported treadmill training improves cardiovascular fitness and walking endurance early after stroke Vol 59 No 4, p 275

##### Clinical Practice Guidelines

- Palliation in aged care Vol 58 No 1, p 63
- Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis Vol 58 No 2, p 128
- Osteoarthritis Vol 58 No 2, p 133
- Hip fracture Vol 58 No 4, p 275
- Osteoporosis Vol 58 No 4, p 275
- Falls in older people Vol 59 No 3, p 213
- Physical activity for older people Vol 59 No 3, p 213

#### Musculoskeletal

##### CAPs

- Targeted physiotherapy treatment for low back pain based on clinical risk can improve clinical and economic outcomes when compared with current best practice Vol 58 No 1, p 57
- A specific exercise program for patients with subacromial impingement syndrome can improve function and reduce the need for surgery Vol 58 No2, p 127
- Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis Vol 58 No 2, p 128
- Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training Vol 58 No 3, p 200

Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272
Exercise programs for patients with cancer improve physical functioning and quality of life	Vol 59 No 1, p 57
Acute low back pain usually resolves quickly but persistent low back pain often persists	Vol 59 No 2, p 127
A physiotherapy telephone assessment and advice service for patients with musculoskeletal problems can improve the process of care while maintaining clinical effectiveness	Vol 59 No 2, p 130
Reconstruction surgery is not always necessary for active young people who rupture their anterior cruciate ligament	Vol 58 No 3, p 209
No difference in functional outcomes between surgery and physiotherapy for symptomatic patients with a meniscal tear and knee osteoarthritis	Vol 59 No 3, p 210

### Clinimetrics

Western Ontario Rotator Cuff Index	Vol 58 No 3, p 201
The patient-rated elbow evaluation (PREE)	Vol 58 No 4, p 274
Pain Free Grip Strength Test	Vol 59 No 1, p 59
STaRT Back Screening Tool	Vol 59 No 2, p 131
The Ten Test for Sensation	Vol 59 No 2, p 132
painDETECT Questionnaire	Vol 59 No 3, p 211
The Work Instability Scale	Vol 59 No 3, p 212

### Clinical Practice Guidelines

Osteoarthritis	Vol 58 No 2, p 133
Hand, hip, and knee osteoarthritis	Vol 58 No 3, p 203
Cervical radiculopathy	Vol 58 No 3, p 203
Hip fracture	Vol 58 No 4, p 275
Osteoporosis	Vol 58 No 4, p 275
Frozen shoulder	Vol 59 No 1, p 61

## Neurology

### CAPs

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia	Vol 58 No 1, p 59
Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No 4, p 269
Tai Chi training is effective in reducing balance impairments and falls in patients with Parkinson's disease	Vol 59 No 1, p 55
Repetitive transcranial magnetic stimulations combined with treadmill training can modulate corticomotor inhibition and improve walking performance in people with Parkinson's disease	Vol 59 No 2, p 128

Repetitive facilitative exercise improves upper limb function in patients with subacute stroke	Vol 59 No 3, p 208
--	--------------------

Body-weight supported treadmill training improves cardiovascular fitness and walking endurance early after stroke	Vol 59 No 4, p 275
---	--------------------

### Clinimetrics

The Dix-Hallpike Test	Vol 58 No 2, p 131
-----------------------	--------------------

### Clinical Practice Guidelines

Cervical radiculopathy	Vol 58 No 3, p 203
Amyotrophic lateral sclerosis (ALS)	Vol 59 No 1, p 61
Paediatric neuromuscular weakness	Vol 59 No 2, p 133

## Orthopaedics

### CAPs

A specific exercise program for patients with subacromial impingement syndrome can improve function and reduce the need for surgery	Vol 58 No2, p 127
Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis	Vol 58 No 2, p 128
Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272
Acute low back pain usually resolves quickly but persistent low back pain often persists	Vol 59 No 2, p 127
Repetitive facilitative exercise improves upper limb function in patients with subacute stroke	Vol 59 No 3, p 208
Reconstruction surgery is not always necessary for active young people who rupture their anterior cruciate ligament	Vol 58 No 3, p 209

### Clinimetrics

The patient-rated elbow evaluation (PREE)	Vol 58 No 4, p 274
Pain Free Grip Strength Test	Vol 59 No 1, p 59
The Ten Test for Sensation	Vol 59 No 2, p 132

### Clinical Practice Guidelines

Osteoarthritis	Vol 58 No 2, p 133
Hand, hip, and knee osteoarthritis	Vol 58 No 3, p 203
Hip fracture	Vol 58 No 4, p 275
Osteoporosis	Vol 58 No 4, p 275
Frozen shoulder	Vol 59 No 1, p 61

## Paediatrics

### CAPs

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia	Vol 58 No 1, p 59
Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy	Vol 58 No 3, p 197

Reconstruction surgery is not always necessary for active young people who rupture their anterior cruciate ligament Vol 58 No 3, p 209

Wearing pedometers in conjunction with daily step goals and incentives can increase physical activity among children Vol 59 No 4, p 273

### Clinical Practice Guidelines

Developmental co-ordination disorder Vol 58 No 2, p 133

Paediatric neuromuscular weakness Vol 59 No 2, p 133

## Sports

### CAPs

Eccentric hamstring muscle training can prevent hamstring injuries in soccer players Vol 58 No 1, p 58

Reconstruction surgery is not always necessary for active young people who rupture their anterior cruciate ligament Vol 58 No 3, p 209

### Clinimetrics

Pain Free Grip Strength Test Vol 59 No 1, p 59

## Other

### CAPs

Positive expiratory pressure prevents more exacerbations than high frequency chest wall oscillation via a vest in people with cystic fibrosis Vol 59 No 4, p 276

### Clinimetrics

Pain Intensity Ratings Vol 58 No 1, p 61

The Dix-Hallpike Test Vol 58 No 2, p 131

Brief Illness Perception Questionnaire (Brief IPQ) Vol 58 No 3, p 202

SPHERE 12 Screening Questionnaire Vol 58 No 4, p 273

Ice-water (cold stress) immersion testing Vol 59 No 4, p 278

# INTERVENTION

## Behaviour Modification

### CAPs

A telephone-delivered behavioural intervention confers a small reduction in body weight in people with Type 2 diabetes Vol 59 No 2, p 133

## Education

### CAPs

Wearing pedometers in conjunction with daily step goals and incentives can increase physical activity among children Vol 59 No 4, p 273

## Fitness Training

### CAPs

Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease Vol 58 No 2, p 129

Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy Vol 58 No 3, p 197

A 12-week exercise program performed during the second trimester does not prevent gestational diabetes in healthy pregnant women Vol 58 No 3, p 198

Interval training confers greater gains than continuous training in people with heart failure Vol 58 No 3, p 199

Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training Vol 58 No 3, p 200

Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility Vol 58 No 4, p 269

Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke Vol 58 No 4, p 271

Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis Vol 58 No 4, p 272

Long-term aerobic exercise maintains peak VO<sub>2</sub>, improves quality of life, and reduces hospitalisations and mortality in patients with heart failure Vol 59 No 1, p 56

Exercise training following lung transplant is now evidence-based practice Vol 59 No 1, p 58

Reconstruction surgery is not always necessary for active young people who rupture their anterior cruciate ligament Vol 58 No 3, p 209

Sun-style T'ai Chi improves walking endurance and health-related quality of life in people with COPD Vol 59 No 4, p 274

Body-weight supported treadmill training improves cardiovascular fitness and walking endurance early after stroke Vol 59 No 4, p 275

### Clinical Practice Guidelines

Physical activity for older people Vol 59 No 3, p 213

## Respiratory Therapy

### CAPs

Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD) Vol 58 No 1, p 60

Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease Vol 58 No 4, p 270

Exercise training following lung transplant is now evidence-based practice Vol 59 No 1, p 58

Telehealth reduces hospital admission rates in patients with COPD	Vol 59 No 2, p 129
Sun-style T'ai Chi improves walking endurance and health-related quality of life in people with COPD	Vol 59 No 4, p 274
Positive expiratory pressure prevents more exacerbations than high frequency chest wall oscillation via a vest in people with cystic fibrosis	Vol 59 No 4, p 276

### Clinical Practice Guidelines

Chronic obstructive lung disease	Vol 59 No 4, p 279
----------------------------------	--------------------

### Strength Training

#### CAPs

Eccentric hamstring muscle training can prevent hamstring injuries in soccer players	Vol 58 No 1, p 58
Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease	Vol 58 No 2, p 129
Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy	Vol 58 No 3, p 197
Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training	Vol 58 No 3, p 200
Repetitive facilitative exercise improves upper limb function in patients with subacute stroke	Vol 59 No 3, p 208
Body-weight supported treadmill training improves cardiovascular fitness and walking endurance early after stroke	Vol 59 No 4, p 275

### Clinical Practice Guidelines

Falls in older people	Vol 59 No 3, p 213
-----------------------	--------------------

### Stretching, Mobilisation, Manipulation, Massage

#### CAPs

Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No 4, p 269
Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke	Vol 58 No 4, p 271
Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272

## PROBLEM

### Difficulty with Sputum Clearance

#### CAPs

Positive expiratory pressure prevents more exacerbations than high frequency chest wall oscillation via a vest in people with cystic fibrosis	Vol 59 No 4, p 276
---	--------------------

### Clinical Practice Guidelines

Chronic obstructive lung disease	Vol 59 No 4, p 279
----------------------------------	--------------------

### Impaired Ventilation

#### CAPs

Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD)	Vol 58 No 1, p 60
Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease	Vol 58 No 4, p 270
Telehealth reduces hospital admission rates in patients with COPD	Vol 59 No 2, p 129
Sun-style T'ai Chi improves walking endurance and health-related quality of life in people with COPD	Vol 59 No 4, p 274
Positive expiratory pressure prevents more exacerbations than high frequency chest wall oscillation via a vest in people with cystic fibrosis	Vol 59 No 4, p 276

### Clinical Practice Guidelines

Chronic obstructive lung disease	Vol 59 No 4, p 279
----------------------------------	--------------------

### Muscle Weakness

#### CAPs

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia	Vol 58 No 1, p 59
Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy	Vol 58 No 3, p 197
Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training	Vol 58 No 3, p 200
Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility	Vol 58 No 4, p 269
Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke	Vol 58 No 4, p 271
Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis	Vol 58 No 4, p 272
Tai Chi training is effective in reducing balance impairments and falls in patients with Parkinson's disease	Vol 59 No 1, p 55
Exercise programs for patients with cancer improve physical functioning and quality of life	Vol 59 No 1, p 57



Repetitive transcranial magnetic stimulations combined with treadmill training can modulate corticomotor inhibition and improve walking performance in people with Parkinson's disease Vol 59 No 2, p 128

Repetitive facilitative exercise improves upper limb function in patients with subacute stroke Vol 59 No 3, p 208

Body-weight supported treadmill training improves cardiovascular fitness and walking endurance early after stroke Vol 59 No 4, p 275

### Clinimetrics

Western Ontario Rotator Cuff Index Vol 58 No 3, p 201

Pain Free Grip Strength Test Vol 59 No 1, p59

### Clinical Practice Guidelines

Frozen shoulder Vol 59 No 1, p 61

Amyotrophic lateral sclerosis (ALS) Vol 59 No 1, p 61

Paediatric neuromuscular weakness Vol 59 No 2, p 133

## Pain

### CAPs

Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis Vol 58 No 4, p 272

Acute low back pain usually resolves quickly but persistent low back pain often persists Vol 59 No 2, p 127

### Clinimetrics

Pain Intensity Ratings Vol 58 No 1, p 61

Active Straight Leg Raise Vol 58 No 2, p 132

Western Ontario Rotator Cuff Index Vol 58 No 3, p 201

The patient-rated elbow evaluation (PREE) Vol 58 No 4, p 274

Pain Free Grip Strength Test Vol 59 No 1, p 59

Chronic Pain Grade Questionnaire Vol 59 No 1, p 60

STaT Back Screening Tool Vol 59 No 2, p 131

painDETECT Questionnaire Vol 59 No 3, p 211

The Work Instability Scale Vol 59 No 3, p 212

### Clinical Practice Guidelines

Osteoarthritis Vol 58 No 2, p 133

Hand, hip, and knee osteoarthritis Vol 58 No 3, p 203

Cervical radiculopathy Vol 58 No 3, p 203

Frozen shoulder Vol 59 No 1, p 61

## Reduced Exercise Tolerance

### CAPs

Interval training confers greater gains than continuous training in people with heart failure Vol 58 No 3, p 199

Group task-specific circuit training for patients discharged home after stroke may be as effective as individualised physiotherapy in improving mobility Vol 58 No4, p 269

Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke Vol 58 No 4, p 271

Long-term aerobic exercise maintains peak VO<sub>2</sub>, improves quality of life, and reduces hospitalisations and mortality in patients with heart failure Vol 59 No 1, p 56

Exercise programs for patients with cancer improve physical functioning and quality of life Vol 59 No 1, p 57

Exercise training following lung transplant is now evidence-based practice Vol 59 No 1, p 58

Wearing pedometers in conjunction with daily step goals and incentives can increase physical activity among children Vol 59 No 4, p 273

Sun-style T'ai Chi improves walking endurance and health-related quality of life in people with COPD Vol 59 No 4, p 274

### Clinimetrics

Rating of Perceived Exertion (RPE) Vol 58 No 1, p 62

### Clinical Practice Guidelines

Physical activity for older people Vol 59 No 3, p 213

## Other

### CAPs

Targeted physiotherapy treatment for low back pain based on clinical risk can improve clinical and economic outcomes when compared with current best practice Vol 58 No 1, p 57

Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis Vol 58 No 2, p 128

Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease Vol 58 No 2, p 129

Exercise training improves cardiovascular fitness in people receiving haemodialysis for chronic renal disease Vol 58 No 2, p 130

Exercise programs for patients with cancer improve physical functioning and quality of life Vol 59 No 1, p 57

Repetitive transcranial magnetic stimulations combined with treadmill training can modulate corticomotor inhibition and improve walking performance in people with Parkinson's disease Vol 59 No 2, p 128

A physiotherapy telephone assessment and advice service for patients with musculoskeletal problems can improve the process of care while maintaining clinical effectiveness Vol 59 No 2, p 130

A telephone-delivered behavioural intervention confers a small reduction in body weight in people with Type 2 diabetes Vol 59 No 2, p 133

Repetitive facilitative exercise improves upper limb function in patients with subacute stroke	Vol 59 No 3, p 208
Reconstruction surgery is not always necessary for active young people who rupture their anterior cruciate ligament	Vol 58 No 3, p 209
No difference in functional outcomes between surgery and physiotherapy for symptomatic patients with a meniscal tear and knee osteoarthritis	Vol 59 No 3, p 210
Wearing pedometers in conjunction with daily step goals and incentives can increase physical activity among children	Vol 59 No 4, p 273
Body-weight supported treadmill training improves cardiovascular fitness and walking endurance early after stroke	Vol 59 No 4, p 275

### Clinimetrics

Rating of Perceived Exertion (RPE)	Vol 58 No 1, p 62
The Dix-Hallpike Test	Vol 58 No 2, p 131
Brief Illness Perception Questionnaire (Brief IPQ)	Vol 58 No 3, p 202
SPHERE 12 Screening Questionnaire	Vol 58 No 4, p 273
Pain Free Grip Strength Test	Vol 59 No 1, p 59
Chronic Pain Grade Questionnaire	Vol 59 No 1, p 60
The Ten Test for Sensation	Vol 59 No 2, p 132
The Work Limitations Questionnaire (WLQ-25)	Vol 58 No 4, p 277
Ice-water (cold stress) immersion testing	Vol 59 No 4, p 278

### Clinical Practice Guidelines

Palliation in aged care	Vol 58 No 1, p 63
Osteoporosis	Vol 58 No 4, p 275
Cardiovascular disease risk	Vol 59 No 2, p 133
Falls in older people	Vol 59 No 3, p 213
Physical activity for older people	Vol 59 No 3, p 213

## BODY PART

### Head & Neck

#### CAPs

Repetitive transcranial magnetic stimulations combined with treadmill training can modulate corticomotor inhibition and improve walking performance in people with Parkinson's disease	Vol 59 No 2, p 128
--	--------------------

#### Clinimetrics

The Dix-Hallpike Test	Vol 58 No 2, p 131
-----------------------	--------------------

### Upper Arm, Shoulder or Shoulder Girdle

#### CAPs

Repetitive facilitative exercise improves upper limb function in patients with subacute stroke	Vol 59 No 3, p 208
--	--------------------

#### Clinimetrics

Western Ontario Rotator Cuff Index	Vol 58 No 3, p 201
The patient-rated elbow evaluation (PREE)	Vol 58 No 4, p 274

### Clinical Practice Guidelines

A specific exercise program for patients with subacromial impingement syndrome can improve function and reduce the need for surgery	Vol 58 No2, p 127
Frozen shoulder	Vol 59 No 1, p 61

### Hand or Wrist

#### CAPs

Bimanual therapy and constraint-induced movement therapy are equally effective in improving hand function in children with congenital hemiplegia	Vol 58 No 1, p 59
--	-------------------

### Clinical Practice Guidelines

Hand, hip, and knee osteoarthritis	Vol 58 No 3, p 203
------------------------------------	--------------------

### Chest

#### CAPs

Action plans and case manager support may hasten recovery of symptoms following an acute exacerbation in patients with chronic obstructive pulmonary disease (COPD)	Vol 58 No 1, p 60
Combined resistance and aerobic training is more effective than aerobic training alone in people with coronary artery disease	Vol 58 No 2, p 129
Neuromuscular electrical stimulation appears to be useful in people with severe chronic obstructive pulmonary disease	Vol 58 No 4, p 270
Long-term aerobic exercise maintains peak VO <sub>2</sub> , improves quality of life, and reduces hospitalisations and mortality in patients with heart failure	Vol 59 No 1, p 56
Telehealth reduces hospital admission rates in patients with COPD	Vol 59 No 2, p 129
Sun-style T'ai Chi improves walking endurance and health-related quality of life in people with COPD	Vol 59 No 4, p 274

### Clinical Practice Guidelines

Paediatric neuromuscular weakness	Vol 59 No 2, p 133
Chronic obstructive lung disease	Vol 59 No 4, p 279

### Thoracic Spine

#### Clinical Practice Guidelines

Cervical radiculopathy	Vol 58 No 3, p 203
------------------------	--------------------

### Lumbar Spine, SIJ or Pelvis

#### CAPs

Targeted physiotherapy treatment for low back pain based on clinical risk can improve clinical and economic outcomes when compared with current best practice	Vol 58 No 1, p 57
---	-------------------

Abdominal muscle feedforward activation in patients with chronic low back pain is largely unaffected by 8 weeks of core stability training Vol 58 No 3, p 200

Acute low back pain usually resolves quickly but persistent low back pain often persists Vol 59 No 2, p 127

### Clinimetrics

Active Straight Leg Raise Vol 58 No 2, p 132

STaRT Back Screening Tool Vol 59 No 2, p 131

painDETECT Questionnaire Vol 59 No 3, p 211

## Perineum or Genito-Urinary System

### Clinimetrics

Exercise training improves cardiovascular fitness in people receiving haemodialysis for chronic renal disease Vol 58 No 2, p 130

## Thigh or Hip

### CAPs

Eccentric hamstring muscle training can prevent hamstring injuries in soccer players Vol 58 No 1, p 58

Multi-modal realignment treatment decreased pain in people with medial tibiofemoral osteoarthritis Vol 58 No 4, p 272

### Clinical Practice Guidelines

Hand, hip, and knee osteoarthritis Vol 58 No 3, p 203

Hip fracture Vol 58 No 4, p 275

## Lower Leg or Knee

### CAPs

Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis Vol 58 No 2, p 128

Reconstruction surgery is not always necessary for active young people who rupture their anterior cruciate ligament Vol 58 No 3, p 209

No difference in functional outcomes between surgery and physiotherapy for symptomatic patients with a meniscal tear and knee osteoarthritis Vol 59 No 3, p 210

### Clinical Practice Guidelines

Osteoarthritis Vol 58 No 2, p 133

Hand, hip, and knee osteoarthritis Vol 58 No 3, p 203

Hip fracture Vol 58 No 4, p 275

## Whole Body/Other

### CAPs

Functional progressive resistance training improves muscle strength but not walking ability in children with cerebral palsy Vol 58 No 3, p 197

A 12-week exercise program performed during the second trimester does not prevent gestational diabetes in healthy pregnant women Vol 58 No 3, p 198

Interval training confers greater gains than continuous training in people with heart failure Vol 58 No 3, p 199

Aerobic treadmill training effectively enhances cardiovascular fitness and gait function for older persons with chronic stroke Vol 58 No 4, p 271

Tai Chi training is effective in reducing balance impairments and falls in patients with Parkinson's disease Vol 59 No 1, p 55

Exercise programs for patients with cancer improve physical functioning and quality of life Vol 59 No 1, p 57

Exercise training following lung transplant is now evidence-based practice Vol 59 No 1, p 58

A physiotherapy telephone assessment and advice service for patients with musculoskeletal problems can improve the process of care while maintaining clinical effectiveness Vol 59 No 2, p 130

A telephone-delivered behavioural intervention confers a small reduction in body weight in people with Type 2 diabetes Vol 59 No 2, p 133

Repetitive facilitative exercise improves upper limb function in patients with subacute stroke Vol 59 No 3, p 208

Wearing pedometers in conjunction with daily step goals and incentives can increase physical activity among children Vol 59 No 4, p 273

Sun-style T'ai Chi improves walking endurance and health-related quality of life in people with COPD Vol 59 No 4, p 274

Body-weight supported treadmill training improves cardiovascular fitness and walking endurance early after stroke Vol 59 No 4, p 275

Positive expiratory pressure prevents more exacerbations than high frequency chest wall oscillation via a vest in people with cystic fibrosis Vol 59 No 4, p 276

### Clinimetrics

Brief Illness Perception Questionnaire (Brief IPQ) Vol 58 No 3, p 202

SPHERE 12 Screening Questionnaire Vol 58 No 4, p 273

Pain Free Grip Strength Test Vol 59 No 1, p 59

Chronic Pain Grade Questionnaire Vol 59 No 1, p 60

The Ten Test for Sensation Vol 59 No 2, p 132

The Work Limitations Questionnaire (WLQ-25) Vol 58 No 4, p 277

### Clinical Practice Guidelines

Palliation in aged care Vol 58 No 1, p 63

Type 2 diabetes Vol 58 No 1, p 63

Developmental co-ordination disorder Vol 58 No 2, p 133

Osteoporosis Vol 58 No 4, p 275

Amyotrophic lateral sclerosis (ALS) Vol 59 No 1, p 61

Cardiovascular disease risk Vol 59 No 2, p 133

Falls in older people Vol 59 No 3, p 213

Physical activity for older people Vol 59 No 3, p 213